

## For better or worse? Changing shift schedules and the risk of work injury among men and women

by Imelda S Wong, PhD,<sup>1</sup> Peter M Smith, MPH, PhD,<sup>1, 2, 3</sup> Cameron A Mustard, PhD,<sup>1, 3</sup> Monique AM Gignac, PhD<sup>1, 3</sup>

Wong IS, Smith PM, Mustard CA, Gignac MAM. For better or worse? Changing shift schedules and the risk of work injury among men and women. *Scand J Work Environ Health*. 2014;40(6):621–630. doi:10.5271/sjweh.3454

**Objectives** The aim of this study was to examine the risk of work injury associated with changes in shift schedules and identify whether work injury risks differ between men and women.

**Methods** Longitudinal panels from the Survey of Labor and Income Dynamics were used to describe work schedule patterns over a 6-year period among a representative sample of Canadian workers (N=19 131). Cox regression was used to estimate the risk of work injury among workers who (i) switched from regular day to nonstandard shifts, (ii) switched from nonstandard to day shifts and (iii) remained in nonstandard shifts, compared with (iv) those who worked regular day shifts only. Gender differences were examined in separate stratified analyses. Adjustments were made for potential respondent and occupational confounders.

**Results** Increased injury risk was observed among those who: switched from day to nonstandard shifts [hazard ratio (HR) 2.60, 95% confidence interval (95% CI) 1.79–3.77], switched from nonstandard to days (HR 2.36, 95% CI 1.62–3.49), and worked nonstandard shifts only (HR 1.44, 95% CI 1.23–1.70). For women, work injury risk was higher among those who switched shifts (days to nonstandard HR 3.10, 95% CI 1.76–5.46; nonstandard to days HR 2.31, 95% CI 1.36–3.91), or worked nonstandard shifts only (HR 1.85, 95% CI 1.44–2.37) compared to day schedules. However, for men the risk of injury was elevated only among those who switched shifts (days to nonstandard HR 2.18, 95% CI 1.35–3.51; nonstandard to days HR 2.38, 95% CI 1.41–3.95). The only significant difference between men and women were among nonstandard shift workers.

**Conclusion** Our results suggest that changing shift types may increase work injury risk among men and women, and that the risk remains increased among women who work nonstandard shifts for a prolonged period of time. This highlights the need for awareness and implementation of health and safety programs when workers initially change shift schedules and on a regular basis to maintain worker health.

**Key terms** Canada; epidemiology; gender difference; nonstandard shift; occupational injury; shift work adjustment; shift work recovery.

It has been estimated that in Canada, similar to other industrialized countries, approximately 30% of the workforce is employed in schedules outside of conventional daytime hours (1). Defined as nonstandard shifts, this type of working schedule includes regular evening or overnight shifts and rotations between day and night shifts (2). With the large number of workers employed in nonstandard shifts (1), potential adverse health outcomes related to these schedules have been of increasing concern. These concerns revolve around findings that biological function-

ing in humans follows a circadian rhythm that promotes activity during daylight hours and recovery when it is dark. Staying awake during normal sleeping hours can lead to circadian desynchronization, changes in normal biological processes, and – over the long term – has been associated with poorer health, including increased risk of chronic disease (3–6). It is believed that work at night also may affect psychosocial and behavioral stressors such as diet and exercise with further deleterious health effects (7). Disruption in duration and quality of sleep attributed

<sup>1</sup> Institute for Work and Health, Toronto, Canada.

<sup>2</sup> School of Public Health & Preventive Medicine, Monash University, Melbourne, Australia.

<sup>3</sup> Dalla Lana School of Public Health, University of Toronto, Canada.

Correspondence to: Imelda S Wong, Institute for Work and Health, Suite 800, 481 University Avenue, Toronto, ON, M5G 2E9, Canada. [E-mail: iwong@iwh.on.ca]

to shift work has been associated with increased fatigue and poorer concentration (8, 9). This combination of problems associated with working night-shifts may have a cumulative effect on overall health and job performance (10) which may contribute to an increased risk of work-related injuries (1, 11–14).

Gender differences in work injury risk among shift workers have also been explored. However, there is limited empirical evidence showing an interaction between shift work and gender that definitively points to shift work as more adverse for either women or men. It has been suggested that night shift work for women may result in greater fatigue and reduced work–life balance and may potential lead to a higher risk of work injury (1, 15). For example, a recent study found that trouble sleeping was associated with work injury among women, but not men, who were working nonstandard shifts (6). In many households, women remain the primary family caregivers (16) and spend more time on household duties, such as childcare, than men (17). The added responsibility of work at night may also contribute to job strain, role overload, and role conflict (18) and ultimately may put women at a higher risk for work injuries (19) although it is not clear whether this effect would remain once gender differences in occupation are taken into consideration. Men are more likely to be employed in physically demanding occupations, which increases their chances of being injured at work (20). However, some studies have found that after accounting for occupational physical demands, work during non-regular daytime schedules may be more harmful for women than men in terms of work-related injuries (1, 21).

In understanding the employment trajectories of workers, it is important to recognize that individuals can change their shift type over time. These changes may be voluntary (eg, to take advantage of new job opportunities or for health and lifestyle preferences or concerns) or involuntary (eg, as a result of organization restructuring). However, there is currently little information on whether changes in a worker's shift type are associated with the risk of work injury over a prolonged period of time. Existing evidence suggests that individuals working regular day shifts have a lesser likelihood than nonstandard shift workers of experiencing work injuries. As such, switching out of nonstandard shifts may provide a "better" working environment in terms of worker safety. The corollary would be that those switching out of day shifts to nonstandard shifts might experience an increased risk of injury as they adjust to being awake and working during normal sleeping hours or deal with variability in work scheduling. In a recent study examining the impact of entering or leaving shift work on sleepiness, results showed that moving from a regular day shift into a nonstandard shift resulted not only in an increased risk of having difficulties with

entering recovery sleep but also with increased risk of falling asleep at work (22). For those switching out of nonstandard shifts, the risk of having difficulties entering recovery sleep was significantly reduced. Similar findings were reported among a study of retired shift workers with sleep quality improving after time away from shift work (23).

The purpose of this longitudinal study is to (i) gain a better understanding of the characteristics of workers who change shift types, (ii) characterize the probability of work injury over time, following a change in shift type, (iii) examine if changing shift type is associated with an increased risk of work injury, and (iv) identify whether work injury risks differ between men and women. This expands upon the existing body of studies, which have focused mainly on the risk of work injury associated with specific shift types such as regular nights or rotating schedules (15) or short-term biological responses of greater fatigue and stress during night shifts and recovery days (24–26). We hypothesize that, compared to workers who work regular day shifts only, those who switch from day to nonstandard shifts will have the highest risk work injury. Workers in nonstandard shifts for a prolonged period of time may have acclimatized to their work schedules and thus are expected to have a lower risk of injury than those who have just recently transitioned from regular day shifts. Workers who change from nonstandard to day shifts should benefit from protective effects of day shifts and are expected to have similar injury risks at those working only day shifts. In keeping with emerging findings of gender differences in previous research, we anticipate that women who are transitioning into, or remaining in, nonstandard shifts will have a higher risk of work-related injury than men.

## Methods

### Data sources and population

This study used data gathered in the Survey of Labor and Income Dynamics (SLID), a household income survey administered by Statistics Canada that collects information regarding an individual's labor market activity and income over six consecutive years. In each survey group, approximately 17 000 households (34 000 respondents) are interviewed annually in January to report on their labor market participation over the previous 12 months. Approximately 80% of respondents provide permission to gather additional information from their federal income tax forms. The remaining 20% of respondents who decline permission are interviewed for a second time in May to provide additional income information

for the previous year (January to December). A stratified, random clustered sampling design provides a representative sample of the non-institutionalized Canadian population. The response rates for the longitudinal panels have decreased with time but still remain high (85% in 1996; 72% in 2010) (27, 28).

Data from three SLID survey groups (1999–2004, 2002–2007, 2005–2010) were used for the study. The study population included all respondents between the ages of 16–69 years who reported working as a paid employee and provided information for all six years of the survey. For respondents who switched shifts more than once during their participation in the SLID, follow-up time was censored to the start of their second shift change. We excluded those with a shift change in their last year of participation in the SLID because injury information post-shift change would not be available in the data.

### Outcome and predictor variables

*Outcome variable: work-related injuries.* Respondents were asked annually whether, over the past 12 months, they received workers' compensation for an injury sustained at work or were absent from work for a period of  $\geq 1$  week that was attributed to a work-related injury or illness. We used the first incidence of workers' compensation or absence as a surrogate measure for incidence of a work injury resulting in lost work days.

*Primary predictor: work schedule.* During each survey, respondents in the SLID were asked to select from eight SLID shift-type categories best describing their work schedule in the past 12 months. We collapsed these categories into two shift work types: (i) regular day shifts and (ii) nonstandard shifts. Using definitions from prior studies (1, 2), we defined "nonstandard shift" workers as those who reported working anything other than a "regular daytime schedule", such as an evening or overnight shift or a rotating day/night shift. A shift change was identified if a respondent reported different shift types in consecutive years during their participation in the SLID.

From this information, four categories were created to describe respondents' work schedules during the 6-year survey period: (i) regular day shifts only (reference); (ii) switched from regular day shifts to nonstandard shifts; (iii) switched from nonstandard shifts to regular day shifts; and (iv) nonstandard shifts only (ie, never worked a regular day shift during survey participation)

*Other independent variables.* We formulated two categories of variables (respondent and occupational), which may confound or modify the relationship between shift work and work injury. Respondent characteristics included age, gender, highest education level, and

household income. Gender was also treated as an effect modifier to account for hypothesized differences in work and home experiences between men and women. We included province of residence to account for potential regional differences in labor market and worker compensation policies. Occupational characteristics (ie, job permanency, usual weekly hours worked, physical demands, and number of employees in the organization) provided information about working conditions that may be associated with the risk for a work injury. Those new to a work situation, such as those starting a job, who work a greater number of hours, or who work in large organizations have been found in previous research to have increased risks of work injury (29–33).

Limited information is available from the SLID regarding occupational demands. However, we included a measure of occupational physical demands in terms of strength requirements typically used while working. This variable was derived from occupational codes following guidelines defined by Human Resources and Skill Development Canada (34) and has been shown to be strongly correlated with work-related injuries (35, 36). Strength requirements were characterized using four categories: (i) limited – tasks that involve handling loads  $< 5$  kg (eg, administrative positions); (ii) light – tasks that involve handling loads of 5–10 kg (eg, shoe repair); (iii) medium – tasks that involve handling loads of 10–20 kg (eg, carpenters); and (iv) heavy – tasks that involve handling loads  $> 20$  kg (eg, road construction). Information regarding respondent and occupational characteristics was taken from the first year of survey enrollment for those who worked day or nonstandard shifts only. For those who switched shift types during the study period, this information was obtained from the first survey following the shift change.

### Statistical analysis

Each respondent who participated in all six years of the SLID survey was identified. For participants with a shift change, the start of their work-injury follow-up (ie, the period in which we examined presence/absence of work injury) was the year the shift change occurred. For individuals who worked day or nonstandard shifts only during the survey period, the work-injury follow-up period began with the first year of their enrollment into the SLID. End of work-injury follow-up was either at the end of the 6-year survey period, when a work injury occurred, or when a second shift change occurred. The total number of annual paid work hours was summed across years of follow-up for each respondent.

Demographic and occupational characteristics were described for each work schedule group. A test of equality for proportions was conducted to compare our study population with the larger workforce population. Chi-

squared tests were used to determine differences in characteristics between shift schedule groups. Incident rates were calculated as the number of incidents over the person-time hours. Incident rate ratios (IRR) were calculated by dividing incident rates for each shift schedule group by rates for daytime only workers. Univariate analyses were completed with Kaplan Meier methods to describe cumulative probability of work injury over time and graphically examine the assumption of proportional hazards. Univariate analyses also showed that work schedule categories violated the assumption of proportional hazards, and therefore this variable was treated as a time-varying variable in our Cox proportional hazard analyses. Log-rank tests determined if differences were significant between groups.

Multivariate analyses were carried out using Cox proportional hazards regression models to assess the risk of work injury over different shift schedules. Adjustments were made for respondent and occupational characteristics, which may confound or modify the risk of work injury. Controlling for year of survey was also included to reflect any policy changes that may have occurred over time and as such, were treated as time-varying covariates. To account for survey non-responses (based on age, gender, geographic region, and income) and the complex sampling design of the SLID, confidence intervals around each estimate were adjusted using a bootstrap technique with 1 000 replicate weights supplied by Statistics Canada (37). Stratified analyses for women and men were completed to examine gender as an effect modifier and to determine risk estimates for each work schedule. Gender differences across independent variables were assessed with a series of interaction terms. Post-hoc pairwise regression comparisons were used to determine significant differences between specific categories of work schedules.

All statistical analyses were completed with Stata version 12 (StataCorp LP, College Station, TX, USA). All P-values  $\alpha=0.05$  were considered statistically significant. Ethical approval for this study was obtained through the University of Toronto, Health Sciences' Ethics Committee.

## Results

In the three SLID survey panels (1999–2004, 2002–2007, and 2005–2010), there were 94 280 respondents aged 16–69 years. Among them, 70 070 (74%) reported being employed at some point during their participation in the survey. We restricted the population to those who provided responses and were employed in all six years of the survey (N=19 898) and excluded those with a shift change in the last year of survey participa-

tion (N=767). Our final population consisted of 19 131 respondents. While this represented only 27% of the population who reported working during some point in their participation in the SLID, both the restricted and full populations were similar in terms of distribution of gender ( $P=0.33$ ) and income ( $P=0.41$ ). However, a larger proportion of our final, restricted population were older ( $P<0.001$ ), had a higher education level ( $P<0.001$ ), and were employed in stable occupational situations (eg, full time employment, permanent positions,  $P<0.001$  for both) (See appendix).

The distribution and characteristics of respondents in each shift schedule group is reported in table 1 (day shifts only 52.0%, day to nonstandard shifts 18.9%, nonstandard to day shifts 17.5%, and nonstandard shifts only 11.5%). Workers who switched from day to nonstandard shifts and from nonstandard to day shifts had more similarities to those who worked nonstandard shifts only than those who worked only regular days (table 1). Specifically, those who worked nonstandard shifts anytime during their participation in the SLID were younger (16–24 years,  $P<0.001$ ) and had lower educational status (less than or equivalent to high school,  $P<0.001$ ) than those employed in regular day shifts only. Working conditions were also similar between those who switched shift types and those employed in nonstandard shifts only: a larger proportion worked <25 hours per week ( $P=0.149$ ) and had jobs requiring medium or heavy strength utilization ( $P=0.174$ ). Switching between shift types appeared to be related to job permanency, as a larger proportion of respondents were employed in temporary positions compared to those who worked day or night shifts only ( $P<0.001$  for both).

The cumulative injury incidence over the SLID's 6-year follow-up period was 8.6%, resulting in an annual injury rate of 1.4% (table 1). This corresponds with national estimates of awarded compensation claims from 2010 (38). While regular daytime-only workers worked more hours than other groups and thus had more opportunity for injury incidence (table 2), the incidence rates and IRR among those who switched shifts were approximately double that of daytime-only workers [day to nonstandard: IRR 2.19, 95% confidence interval (95% CI) 1.91–2.50; nonstandard to day: IRR 1.90, 95% CI 1.66–2.16; nonstandard shifts only: IRR 1.99, 95% CI 1.78–2.22].

Kaplan Meier analyses revealed that the cumulative probability of injury increased at different rates across categories of work schedules for all workers (figure 1). Results of log-rank tests indicated that probability of injury among those working day shifts only was significantly lower than all other work schedules ( $P<0.001$ ).

Unadjusted injury hazard ratios (HR) (not shown) resulted in significantly higher risks among all categories of work schedules compared to those who worked




**Table 1.** Respondent and occupational characteristics.

	Regular day shifts only (N=9948)		Regular day to nonstandard shifts (N=3622)		Nonstandard to regular day shifts (N=3344)		Nonstandard shifts only (N=2217)	
	N	%	N	%	N	%	N	%
Work injury								
Yes	1166	11.7	272	7.5	290	8.7	467	21.0
No	8782	88.3	3350	92.5	3054	91.3	1750	78.9
Sex								
Male	4779	48.0	1853	51.2	1677	50.1	1140	51.4
Female	5169	52.0	1769	48.8	1667	49.9	1077	48.6
Age group (years)								
16–24	503	5.1	513	14.2	827	24.7	276	12.4
25–34	2005	20.2	719	19.9	694	20.8	445	20.1
35–44	3598	36.2	959	26.5	815	24.4	723	32.6
45–54	3096	31.1	959	26.5	694	20.8	587	26.5
≥55	746	7.5	472	13.0	314	9.4	186	8.4
Education								
High school or less	3512	35.6	1685	46.9	1599	48.2	1032	47.0
Non-university certificate	3504	35.5	1268	35.3	1184	35.7	924	42.1
University	2846	28.9	637	17.7	536	16.1	240	10.9
Annual household income (\$)								
<20 000	918	9.2	540	14.9	468	14.0	270	12.2
20 000–40 000	5111	51.4	1764	48.7	1614	48.3	1236	55.8
40 000–60 000	2917	29.3	936	25.8	906	27.1	564	25.4
>60 000	1002	10.1	382	10.5	356	10.6	147	6.6
Weekly work hours								
1–25	643	6.8	826	23.9	572	18.0	428	20.3
26–35	2045	21.5	623	18.1	653	20.6	349	16.5
36–40	5604	59.0	1222	35.4	1319	41.5	892	42.3
≥41	1208	12.7	778	22.6	632	19.9	442	20.9
Permanent job								
Temporary	883	8.9	878	24.3	783	23.4	279	12.6
Permanent	9057	91.1	2737	75.7	2558	76.6	1935	87.4
Strength requirement								
Limited	6173	64.3	1545	43.4	1400	42.7	710	32.6
Light	1467	15.3	886	24.9	861	26.3	715	32.8
Medium	1451	15.1	859	24.1	749	22.8	510	23.4
Heavy	508	5.3	271	7.6	269	8.2	246	11.3
Number of employees								
<20	1854	18.6	781	21.6	757	22.6	237	10.7
20–99	1684	16.9	591	16.3	537	16.1	274	12.4
100–499	1373	13.8	527	14.5	469	14.0	327	14.7
≥500	5037	50.6	1723	47.6	1581	47.3	1379	62.2

day shifts only (day to nonstandard shifts HR 2.47, 95% CI 2.06–2.96; nonstandard to day shifts HR 1.82, 95% CI 1.52–2.17; and nonstandard shifts only HR 1.89, 95% CI 1.64–2.17). In the final model, the risk of injury remained significant after adjustment for respondent and occupational characteristics (table 3). Increased injury risk was observed among those who: switched from day to nonstandard shifts (HR 2.60, 95% CI 1.79–3.77), switched from nonstandard shifts to days (HR 2.36, 95% CI 1.62–3.49), and who remained in nonstandard shifts during the study period (HR 1.44, 95% CI 1.23–1.70). There was no difference in risk between workers who switched shift types ( $P=0.14$ ), and those who worked nonstandard shifts only ( $P=0.41$ ). Significantly increased injury risks remained among women and those with less than university-level education, lower incomes, in physically demanding occupations, in non-permanent

positions, or working <35 hours per week. We also found that those employed in establishments with <20 employees were less likely to be awarded workers' compensation or have a work-injury absence for >7 days.

In the gender-stratified analysis (table 3), the only significant difference between men and women were among those working nonstandard shifts only ( $P<0.001$ ). Within-gender results showed that the risk of work injury was slightly higher for women who switched shifts (days to nonstandard HR 3.10, 95% CI 1.76–5.46; nonstandard to days HR 2.31, 95% CI 1.36–3.91), or worked only nonstandard shifts (HR 1.85, 95% CI 1.44–2.37). Risks were not significantly different between those who switched shift types ( $P=0.23$ ), but both groups had a significantly higher risk of work injury compared with women working nonstandard shifts only ( $P<0.001$ ). Among men, the risk of injury remained higher among



**Figure 1.** Kaplan Meier hazard curve for cumulative probability of work-related injury by work schedule.

**Table 2.** Incidence rates (IR) and incidence rate ratios (IRR) by shift schedule [95% CI=95% confidence interval]

	Person hours of exposure (million person-hrs)	IR (million person-hours)	IRR	95% CI
Day shifts only	104.5	11.2	Reference	
Day to nonstandard shifts	11.1	24.4	2.19	1.91–2.50
Nonstandard to day shifts	13.7	21.2	1.90	1.66–2.16
Nonstandard shifts only	21.0	22.2	1.99	1.78–2.22

those who switched shifts (days to nonstandard HR 2.18, 95% CI 1.35–3.51; nonstandard to days HR 2.38, 95% CI 1.41–3.95), with no difference between both these groups ( $P=0.73$ ). However, the risk among male night-shift-only workers was attenuated and no longer significant (HR 1.29, 95% CI 0.99–1.61).

## Discussion

Existing evidence has focused primarily on the risk of work injury or short-term recovery associated with nonstandard shifts. This study fills a gap in our understanding of shift work by examining work injuries among those workers who switch shift schedules. Previous studies have shown that those employed in regular daytime schedules are less likely to be injured on the job, suggesting that regular daytime workers may have a “better” work schedule in terms of prevention of work

injuries. As such, we expected that respondents switching from nonstandard to daytime-only shifts would benefit from this “better” schedule with respect to work injuries. However, the results of this study shows that work injury risk associated with changing shift schedules is more complex than previously assumed.

This study extended prior research to examine (i) the probability of sustaining a work-related injury following a change in shift type, (ii) whether work injury risk was associated with changes in shift type, and (iii) whether the pattern of risks were different between women and men. We found that personal and occupational characteristics of those whose shifts changed were often similar to those who worked nonstandard shifts compared to those who worked only regular days. Specifically, workers who changed shifts were generally younger and had lower education and income levels. In prior studies, these same factors have been associated with an increased risk of work injury (39). Moreover, occupational characteristics that were reported among those who changed shift types, namely physically demanding jobs and non-permanent positions also have been identified as factors that are associated with increases in work injury risk (30, 40). Of interest in this research is that even after adjustments for these factors, increased risk of work injury was still present among those who switched shifts. Future studies are needed to replicate these findings and examine other occupational factors (eg, specific job tasks) that may play a role along with changing shift types and the increase of work injury risk. This information could be used to help develop occupational health and safety programs to maintain worker health.

**Table 3.** Injury hazard ratios (HR) for principal independent variables for the fully adjusted model and stratified by gender. [95% CI=95% confidence interval]

	Fully adjusted model <sup>a</sup>		Stratified by gender <sup>b</sup>				Difference between men and women
	All workers		Men		Women		
	HR	95% CI	HR	95% CI	HR	95% CI	P <sub>interaction</sub>
Work schedule							
Day shifts only	reference		reference		reference		
Day to nonstandard shifts	2.60 <sup>c</sup>	1.79–3.77	2.18 <sup>c</sup>	1.35–3.51	3.10 <sup>c</sup>	1.76–5.46	0.79
Nonstandard to day shifts	2.36 <sup>c</sup>	1.62–3.49	2.38 <sup>c</sup>	1.41–3.95	2.31 <sup>c</sup>	1.36–3.91	0.18
Nonstandard shifts only	1.44 <sup>c</sup>	1.23–1.70	1.29	0.99–1.61	1.85 <sup>c</sup>	1.44–2.37	P<0.001
Age (years)							
16–24	1.00	0.80–1.26	0.88	0.65–1.20	1.05	0.75–1.47	0.44
25–34	1.04	0.87–1.23	1.07	0.85–1.34	0.95	0.73–1.23	0.56
35–44	reference		reference		reference		
45–54	1.09	0.94–1.27	1.14	0.94–1.39	1.07	0.85–1.34	0.53
≥55	1.14	0.92–1.42	1.15	0.85–1.55	1.14	0.83–1.58	0.38
Sex							
Male	reference						
Female	1.16 <sup>c</sup>	1.01–1.33					
Education							
High school or less	1.75 <sup>c</sup>	1.41–2.15	2.75 <sup>c</sup>	1.94–3.89	1.26	0.96–1.67	P<0.001
Non-university certificate	1.63 <sup>c</sup>	1.32–2.00	2.43 <sup>c</sup>	1.73–3.42	1.25	0.95–1.64	P<0.001
University	reference		reference		reference		
Household income (\$)							
<20 000	1.48 <sup>c</sup>	1.08–2.01	1.33	0.87–2.05	1.75 <sup>c</sup>	1.12–2.74	0.85
20 000–40 000	1.71 <sup>c</sup>	1.32–2.22	1.68 <sup>c</sup>	1.16–2.42	1.81 <sup>c</sup>	1.26–2.61	0.60
40 000–60 000	1.19	0.91–1.55	1.16	0.80–1.69	1.23	0.85–1.79	0.86
≥60 000	reference		reference		reference		
Job permanency							
Temporary	reference		reference		reference		
Permanent	0.81 <sup>c</sup>	0.67–0.99	0.70 <sup>c</sup>	0.54–0.91	1.01	0.76–1.36	0.03
Weekly working hours							
1–25	1.35 <sup>c</sup>	1.08–1.69	1.52 <sup>c</sup>	1.01–2.30	1.34 <sup>c</sup>	1.02–1.75	0.41
26–35	1.30 <sup>c</sup>	1.11–1.53	1.49 <sup>c</sup>	1.16–1.92	1.18	0.96–1.45	0.38
36–40	reference		reference		reference		
≥41	0.88	0.74–1.04	0.81 <sup>c</sup>	0.66–0.99	1.06	0.76–1.48	0.62
Strength requirement							
Limited	reference		reference		reference		
Light	1.78 <sup>c</sup>	1.50–2.10	1.99 <sup>c</sup>	1.54–2.56	1.66 <sup>c</sup>	1.31–2.09	0.04
Medium	2.36 <sup>c</sup>	2.01–2.77	2.58 <sup>c</sup>	2.04–3.25	1.92 <sup>c</sup>	1.50–2.45	0.01
Heavy	2.27 <sup>c</sup>	1.83–2.82	2.49 <sup>c</sup>	1.92–3.25	1.41	0.76–2.63	0.03
Number of employees							
<20	0.73 <sup>c</sup>	0.61–0.87	0.90	0.72–1.14	0.50 <sup>c</sup>	0.38–0.65	P<0.001
20–99	0.90	0.77–1.06	1.01	0.81–1.25	0.79	0.62–1.00	0.06
100–499	1.01	0.85–1.19	0.83	0.66–1.05	1.25	0.99–1.58	0.02
≥500	reference		reference		reference		

<sup>a</sup> Adjusted for age, sex, household income, job permanency, weekly working hours, occupational strength requirements, number of employees.<sup>b</sup> Fully adjusted model stratified by gender.<sup>c</sup> P<0.05.

Our results supported the hypothesis that those who switched from regular days to nonstandard shifts would have the highest injury risk. It may be that those new to working nonstandard shifts experience greater fatigue or other difficulties that may increase their risk for work injury. Of concern, however, were findings that those who switched from nonstandard to day shifts had a similar injury risk to those switching in the opposite direction. The explanation for this finding is unclear. Our data did not provide information on the reasons for a change in shift type. It may be that a change in shift was associated with more hazardous work environments,

thereby increasing the risk for injury, or poorer health precipitated a shift change and continued to exert a potential impact on increasing the risk of a work injury. Further investigation is needed to understand lifestyle and occupational factors that may have a unique impact on some types of shift changes more than others or that might have similar impacts on a range of work scheduling transitions and the risk of work injury. Finally, those who worked nonstandard shifts only were more likely to be injured at work compared with those who remained in day shifts during the study period. This is similar to findings from previous studies (1, 15). As expected,

individuals who worked nonstandard shifts only were less likely to be injured than those who switched shift types. Additional research is needed, but it may be that those who remain in nonstandard shifts for a prolonged period of time are able to adjust to the potentially adverse conditions.

Stratified analyses revealed gender differences for the risk of work injury for respondents who remained in nonstandard shifts, but no differences between men and women who changed shift schedule (in either direction). Results showed that women had a significantly higher injury risk than men when remaining in nonstandard shifts. This finding adds to evidence from previous studies. It may be linked to differences in fatigue or other lifestyle demands (1, 19) or to differences in health and other occupational exposures. Recent studies have shown that women generally report poorer health, restrictions on performing daily activities and more chronic conditions than men (41). Gender differences across some industries, such as healthcare and manufacturing, may also result in different hazard exposures among men and women and influence the risk of work injury. Additional research is needed to pursue whether differences in injury risk are explained by gender, lifestyle factors (eg, care-giving and household responsibilities), the nature of the work experiences of women and men or other variables such as access to compensation for injury. Important to note is that there were no gender differences associated with switching shifts suggesting that the impact of changes in scheduling is equally disruptive and associated with elevated risks of injury for both women and men.

Within-gender analyses revealed that for men, increased injury risk was significant among those who recently switched from days to nonstandard shifts, but not among those who remained in nonstandard shifts. These findings suggest that for men, there may be an initial period of adjustment to working at night that increases their risk for injury, but over time the risks are diminished. For women, those who worked nonstandard shifts only were less likely to report a work injury as women who had recently started nonstandard shifts, but the risk of injury was still almost twice that of women who worked regular day shifts. Both sets of findings speak to the importance of implementing occupational health and safety policies and programs early after changes in shifts. They also point to the potential importance of greater awareness and health and safety action on an ongoing basis. Additional research is needed to examine whether this may be especially true among women.

This study has several limitations that should be addressed in future research. Our secondary analysis of administrative data did not allow us to explore what circumstances may have influenced a respondent to

switch shift types. For example, switches to nonstandard shifts may have been voluntary or related to pay incentives, because of organizational changes or health factors. These same factors may have also influenced the risk of subsequent work-related injury. The strict inclusion criteria requiring responses from all six years of participation in the SLID to form the study population may have also influenced findings. A sensitivity analysis with the entire workforce population, regardless of response completeness, was not possible with these data. However, a comparison of demographic characteristics suggests that risks for work injury may have been underestimated given that the characteristics of our study population (ie, higher education and stable occupational situations) may have resulted in less hazardous working conditions than the general working population. Finally, the use of reports of workers' compensation and work injury absences for >1 week is a more stringent proxy for work injury than most studies and may have excluded many types of acute injuries. By identifying work injuries with these criteria we have highlighted more severe injuries or claims. Some studies have shown that less than half of all workplace injuries are reported (42) and that the nature of the injury is related to absence duration, with acute traumatic injuries resulting in less time away from work than sprain and strain injuries (43). This may mean that our results underestimated the true risk for workplace injuries related to shift work and schedule change.

Despite some limitations, the longitudinal design of the survey allowed us to follow respondents over a 6-year period to examine the subsequent risk of work-related injury among different work schedules. The annual response to the survey also enabled the identification of changes in work schedules using a large representative sample of the Canadian labor force, with information provided from federal income tax forms for 96% of respondents. The findings highlight concerns about the risk of injury, not only among both men and women who switch from day to nonstandard shifts, but also about injury risks remaining elevated particularly among women who work nonstandard shifts for a prolonged period of time. In addition, the benefits of switching from nonstandard to day shifts may not reduce injury risk among men or women. These results have implications for health and safety programs and the need to develop additional strategies to help reduce injury risk among those who change shifts, as well as the need to provide awareness, training, and risk management efforts to reduce injuries on an ongoing basis.



## Acknowledgments

The authors would like to acknowledge that funding for I Wong was provided by the Mustard Post-Doctoral Fellowship from the Institute for Work and Health; P Smith is supported by a Discovery Early Career Researcher Award from the Australian Research Council. The Institute for Work & Health operates with the support of the Province of Ontario. The views expressed in this document are those of the authors and do not necessarily reflect those of the Province of Ontario.

## References

- Wong IS, McLeod CB, Demers PA. Shift work trends and risk of work injury among Canadian workers. *Scand J Work Environ Health*. 2011 Jan;37(1):54–61. <http://dx.doi.org/10.5271/sjweh.3124>.
- Dembe AE, Delbos R, Erickson JB, Banks SM. Associations between employees' work schedules and the vocational consequences of workplace injuries. *J Occup Rehab*. 2007 Dec;17(4):641–51. <http://dx.doi.org/10.1007/s10926-007-9098-8>.
- Davis S, Mirick DK. Circadian disruption, shift work and the risk of cancer: a summary of the evidence and studies in Seattle. *Cancer Causes Control*. 2006 May;17(4):539–45. <http://dx.doi.org/10.1007/s10552-005-9010-9>.
- Erren TC, Morfeld P, Stork J, Knauth P, Von Mülmann MJ, Breitstadt R, et al. Shift work, chronodisruption and cancer?—the IARC 2007 challenge for research and prevention and 10 theses from the Cologne Colloquium 2008. *Scand J Work Environ Health*. 2009 Jan;35(1):74–9. <http://dx.doi.org/10.5271/sjweh.1303>.
- Scheer F, Hilton M, Mantzoros C, Shea S. Adverse metabolic and cardiovascular consequences of circadian misalignment. *Proceedings of the National Academy of Sciences of the United States of America*. 2009 Mar 17;106(11):4453–8. <http://dx.doi.org/10.1073/pnas.0808180106>.
- Kling RN, McLeod CB, Koehoorn M. Sleep problems and workplace injuries in Canada. *Sleep*. 2010 May;33(5):611–8.
- Puttonen S, Harma M, Hublin C. Shift work and cardiovascular disease - pathways from circadian stress to morbidity. *Scand J Work Environ Health*. 2010;36(2):96–108. <http://dx.doi.org/10.5271/sjweh.2894>.
- Reid K, Dawson D. Comparing performance on a simulated 12 hour shift rotation in young and older subjects. *Occup Environ Med*. 2001 Jan;58(1):58–62. <http://dx.doi.org/10.1136/oem.58.1.58>.
- Monk TH, Buysse DJ, Billy BD, Fletcher ME, Kennedy KS. Polysomnographic Sleep and Circadian Temperature Rhythms as a Function of Prior Shift Work Exposure in Retired Seniors. *Healthy Aging Clin Care Elder*. 2013;2013(5):9–19. <http://dx.doi.org/10.4137/HACCE.S11528>.
- Akerstedt T, Folkard S, Portin C. Predictions from the three-process model of alertness. *Aviat Space Environ Med*. 2004;75(3 Sup):A75–83.
- Dembe AE, Erickson JB, Delbos RG, Banks SM. Nonstandard shift schedules and the risk of job-related injuries. *Scand J Work Environ Health*. 2006;32(3):232–40. <http://dx.doi.org/10.5271/sjweh.1004>.
- Dembe AE, Delbos R, Erickson JB. The effect of occupation and industry on the injury risks from demanding work schedules. *J Occup Environ Med*. 2008 Oct;50(10):1185–94. <http://dx.doi.org/10.1097/JOM.0b013e31817e7bf2>.
- Horwitz IB, McCall BP. The impact of shift work on the risk and severity of injuries for hospital employees: an analysis using Oregon workers' compensation data. *Occ Med*. 2004 Dec;54(8):556–63. <http://dx.doi.org/10.1093/occmed/kqh093>.
- De Castro AB, Fujishiro K, Rue T, Tagalog E, Samaco-Paquiz LPG, Gee GC. Associations between work schedule characteristics and occupational injury and illness. *Int Nurs Rev*. 2010;57(2):188–94. <http://dx.doi.org/10.1111/j.1466-7657.2009.00793.x>.
- Wagstaff AS, Sigstad Lie J-A. Shift and night work and long working hours – a systematic review of safety implications. *Scand J Work Environ Health*. 2011 May 3;37(3):173–85. <http://dx.doi.org/10.5271/sjweh.3146>.
- Higgins C, Duxbury L, Lee C. Balancing Work and Family: A Study of the Canadian Private Sector Employees. National Centre for Management Research and Development, Western Business School, the University of Western Ontario. London; 1992.
- Berntsson L, Lundberg U, Krantz G. Gender differences in work-home interplay and symptom perception among Swedish white-collar employees. *J Epidemiol Community Health*. 2006 Dec;60(12):1070–6. <http://dx.doi.org/10.1136/jech.2005.042192>.
- Gignac M, Backman C, Kaptein S, Lacaille D, Beaton D, Hofstetter C, Badley EM. Tension at the borders: perceptions of role overload, conflict, strain and facilitation in work, family and health roles among employed individuals with arthritis. *Rheumatology (Oxford, England)*. 2012 Feb;51(2):324–32. <http://dx.doi.org/10.1093/rheumatology/ker317>.
- Smith PM, Ibrahim-Dost J, Keegel T, MacFarlane E. Gender differences in the relationship between shift work and work injury: examining the influence of dependent children. *J Occup Environ Med*. 2013;55(8):932–936. <http://dx.doi.org/10.1097/JOM.0b013e31829178e1>.
- Smith P, Bielecky A, Koehoorn M, Beaton D, Ibrahim S, Mustard C, Saunders R, Scott-Marshall H. Are Age-Related Differences in the Consequence of Work Injury Greater When Occupational Physical Demands are High? *Am J Ind Med*. 2014;57(4):438–44. <http://dx.doi.org/10.1002/ajim.22303>.
- Mustard C, Chambers A, McLeod C, Bielecky A, Smith P. Work injury risk by time of day in two population-based data sources. *Occup Environ Med*. 2013 Jan;70(1):49–56. <http://dx.doi.org/10.1136/oemed-2012-100920>.

22. Akerstedt T, Nordin M, Alfredsson L, Westerholm P, Kecklund G. Sleep and Sleepiness: Impact of entering or leaving shiftwork - a prospective study. *Chronobiol Int*. 2010 Jul;27(5):987–96. <http://dx.doi.org/10.3109/07420528.2010.489423>.
23. Guo Y, Liu Y, Huang X, Rong Y, He M, Wang Y, et al. The effects of shift work on sleeping quality, hypertension and diabetes in retired workers. *PloS one*. 2013 Jan;8(8):e71107. <http://dx.doi.org/10.1371/journal.pone.0071107>.
24. Lindholm H, Sinisalo J, Ahlberg J, Hirvonen A, Hublin C, Partinen M, et al. Attenuation of vagal recovery during sleep and reduction of cortisol/melatonin ratio in late afternoon associate with prolonged daytime sleepiness among media workers with irregular shift work. *Am J Ind Med*. 2012;55(7):643–9. <http://dx.doi.org/10.1002/ajim.22042>.
25. Wehrens S, Hampton S, Skene D. Heart rate variability and endothelial function after sleep deprivation and recovery sleep among male shift and non-shift workers. *Scand J Work Environ Health*. 2012;38(2):171–81. <http://dx.doi.org/10.5271/sjweh.3197>.
26. Wong IS, Ostry AS, Demers PA, Davies HW. Job strain and shift work influences on biomarkers and subclinical heart disease indicators: a pilot study. *J Occup Environ Hyg*. 2012 Jan;9(8):467–77. <http://dx.doi.org/10.1080/15459624.2012.693831>.
27. Bastien J, Cauchon C, Jocelyn W, Ann Lim M. Data Quality for the 2008 Survey of Labour and Income Dynamics [Internet]. 2009 [cited 2014 Jun 22]. Available from: [http://www23.statcan.gc.ca/imdb-bmdi/document/3889\\_D13\\_T2\\_V2-eng.pdf](http://www23.statcan.gc.ca/imdb-bmdi/document/3889_D13_T2_V2-eng.pdf).
28. Statistics Canada. Data Quality for the 2006 Survey of Labour and Income Dynamics (SLID) [Internet]. 2009 [cited 2014 Jun 22]. Available from: <http://www.statcan.gc.ca/pub/75f0002m/2008005/section5-eng.htm>.
29. Breslin FC, Tompa E, Zhao R, Pole JD, Amick Iii BC, Smith PM, Hogg-Johnson S. The relationship between job tenure and work disability absence among adults: a prospective study. *Accid Anal Prev*. 2008 Jan;40(1):368–75. <http://dx.doi.org/10.1016/j.aap.2007.07.007>.
30. Quinlan M, Mayhew C. Precarious employment and workers' compensation. *Int J Law Psychiatry*. 1999;22(5-6):491–520. [http://dx.doi.org/10.1016/S0160-2527\(99\)00023-0](http://dx.doi.org/10.1016/S0160-2527(99)00023-0).
31. Arlinghaus A, Lombardi DA, Willetts JL, Folkard S, Christiani DC. A Structural Equation Modeling Approach to Fatigue-related Risk Factors for Occupational Injury. *Am J Epidemiol*. 2012 Oct 1;176(7):597–607. <http://dx.doi.org/10.1093/aje/kws219>.
32. Lombardi D, Folkard S, Willetts J, Smith G. Daily sleep, weekly working hours, and risk of work-related injury: US National Health Interview Survey (2004-2008). *Chronobiol Int*. 2010 Jul;27(5):1013–30. <http://dx.doi.org/10.3109/07420528.2010.489466>.
33. Oleinick A, Gluck J V, Guire KE. Establishment size and risk of occupational injury. *American journal of industrial medicine*. 1995 Jul;28(1):1–21. <http://dx.doi.org/10.1002/ajim.4700280102>.
34. Human Resources and Skills Development Canada. National Occupational Classification Career Handbook, Second Edition. Physical Activities and Environmental Conditions. (Report number LM-410-05-03E) [Internet]. 2003 [cited 2013 Feb 18]. Available from: <http://www5.hrsdc.gc.ca/noC/English/CH/2001/physicalActivities.aspx>.
35. Smith P, Bielecky A, Mustard C. The relationship between chronic conditions and work-related injuries and repetitive strain injuries in Canada. *J Occup Environ Med*. 2012 Jul;54(7):841–6. <http://dx.doi.org/10.1097/JOM.0b013e31824e11f7>.
36. Smith P, Bielecky A, Mustard C, Beaton D, Hogg-Johnson S, Ibrahim S, Koehoorn M, McLeod C, Saunders R, Scott-Marshall H. The Relationship between Age and Work Injury in British Columbia : Examining Differences across Time and Nature of Injury. *J Occup Health*. 2013;55(2):98–107. <http://dx.doi.org/10.1539/joh.12-0219-OA>.
37. Yeo D, Mantel H, Liu T. Bootstrap variance estimation for the national population health survey. In the 1999 Proceedings of the Survey Research Methods Section, Americal Statistical Association Conference. Baltimore; 1999 p778–83.
38. Association of Workers' Compensation Boards of Canada. Key Statistical Measures [Internet]. 2010 [cited 2014 Feb 25]. Available from: <https://aoc.awcbc.org/KsmReporting/ReportDataConfig>.
39. Piha K, Laaksonen M, Martikainen P, Rahkonen O, Lahelma E. Socio-economic and occupational determinants of work injury absence. *Eur J Public Health*. 2013 Dec 3;23(4):1–5. <http://dx.doi.org/10.1093/eurpub/cks162>.
40. Smith PM, Kosny AA, Mustard CA. Differences in Access to Wage Replacement Benefits for Absences Due to Work-Related Injury or Illness in Canada. *Am J Ind Med*. 2009;52(4):341–9. <http://dx.doi.org/10.1002/ajim.20683>.
41. Oksuzyan A, Crimmins E, Saito Y, O'Rand A, Vaupel JW, Christensen K. Cross-national comparison of sex differences in health and mortality in Denmark, Japan and the US. *Eur J Epidemiol*. 2010 Jul;25(7):471–80. <http://dx.doi.org/10.1007/s10654-010-9460-6>.
42. Shannon HS, Lowe GS. How Many Injured Workers Do Not File Claims for Workers' Compensation Benefits ? *Am J Ind Med*. 2002;42(6):467–73. <http://dx.doi.org/10.1002/ajim.10142>.
43. Health and Safety Executive. Self-reported work-related illness (SWI) and workplace injuries: Results from the Labour Force Survey (LFS) [Internet]. 2012 [cited 2014 Jun 23]. Available from: <http://www.hse.gov.uk/STATISTICS/lfs/index.htm#msd>.

Received for publication: 12 April 2014